

OPTIMAL HÖLDER EXPONENT FOR THE SLE PATH

FREDRIK JOHANSSON AND GREGORY F. LAWLER

ABSTRACT. We prove an upper bound on the optimal Hölder exponent for the chordal SLE path parameterized by capacity and thereby establish the optimal exponent as conjectured by J. Lind. We also give a new proof of the lower bound. Our proofs are based on the sharp estimates of moments of the derivative of the inverse map. In particular, we improve an estimate of the second author.

1. INTRODUCTION

The Schramm-Loewner evolution, or $SLE(\kappa)$, is a one-parameter family of random fractal curves that was introduced by O. Schramm in [Sch00] as a candidate for the scaling limit of the loop-erased random walk. Since then, SLE has been shown to describe the scaling limits of a number of discrete models from statistical physics and to provide tools for their rigorous understanding. The properties of SLE curves have been studied by a number of authors. For instance, S. Rohde and Schramm proved in [RS05] the existence and continuity of the path and gave an upper bound on the Hausdorff dimension. V. Beffara [Bef08] proved a lower bound on Hausdorff dimension and thus showed that the dimension is almost surely the minimum of $1 + \kappa/8$ and 2. J. Lind [Lin08] improved the estimates by Rohde and Schramm and proved that the $SLE(\kappa)$ path is almost surely Hölder continuous. She conjectured that the Hölder exponent she obtained is the optimal one, that is, that the exponent is the largest possible. In this paper we prove this conjecture. More precisely, we prove the following theorem. Let $\kappa \geq 0$ and set

$$\alpha_* = \alpha_*(\kappa) = 1 - \frac{\kappa}{24 + 2\kappa - 8\sqrt{8 + \kappa}}, \quad \alpha_0 = \min \left\{ \frac{1}{2}, \alpha_* \right\}.$$

Theorem 1.1. *Let $\gamma(t)$ be the chordal $SLE(\kappa)$ path parameterized by half-plane capacity. With probability one the following holds.*

Johansson is supported by grant KAW 2005.0098 from the Knut and Alice Wallenberg Foundation.

Lawler is supported by National Science Foundation grant DMS-0734151.




FIGURE 1. Optimal Hölder exponent as a function of κ .

- $\gamma(t)$, $t \in [0, 1]$, is Hölder continuous of order α for $\alpha < \alpha_0$ and is not Hölder continuous of order α for $\alpha > \alpha_0$.
- For every $0 < \epsilon < 1$, $\gamma(t)$, $t \in [\epsilon, 1]$, is Hölder continuous of order α for $\alpha < \alpha_*$ and is not Hölder continuous of order α for $\alpha > \alpha_*$.

As mentioned, the lower bound on the optimal exponent was proved in [Lin08]. The upper bound is new and we also give a new proof of the lower bound. The phase transition of $\alpha_0(\kappa)$ at $\kappa = 1$ is due to the geometry of the path at the base; roughly, in the capacity parameterization the path is Hölder-1/2 at $t = 0$ for all $\kappa \geq 0$.

The main tool needed is sharp estimates for the moments of the derivative close to the preimage of the tip of the growing curve. We will use the work in [Law09] and also [JL09], but we will have to extend one of these results in this paper. We will use these results as a “black box”, more or less, in the main part of the paper and then give a derivation in the final section, building on the argument in [Law09]. To be more specific, we need to control the growth of the derivative as the preimage of the tip is approached radially. To get the lower bound on the exponent estimates on the derivative from above are needed, and to get the upper bound on the optimal exponent, one has to control second moments and time correlations.

Our paper is organized as follows. In Section 2 we give definitions and discuss some well-known results. In Section 3 we prove a number of results about Loewner chains that are not particular to SLE, but rather to Loewner chains driven by functions which are weakly Hölder-1/2,

see (13) for a definition. Notably, we give estimates on the modulus of continuity of the curve in terms of the radial growth of the derivative. In the Section 4 we state the basic moment estimates and use these together with the results from Section 3 to prove Theorem 1.1. Finally, Section 5 proves the estimates on the derivative. Here we review without proof the main results from [Law09] that we need, and then establish the new estimates.

2. PRELIMINARIES

Let U_t , $t \geq 0$ be a continuous real-valued function. We shall consider chordal Loewner chains (g_t) , that is, solutions to the chordal Loewner equation

$$\partial_t g_t = \frac{a}{g_t - U_t}, \quad t > 0, \quad g_0(z) = z, \quad (1)$$

for $a > 0$ fixed. We define the associated continuously growing hull

$$K_t = \{z \in \mathbb{H} : \tau(z) \leq t\},$$

where $\tau(z)$ is the blow-up time of (1). For each $t > 0$ the function $z \mapsto g_t(z)$ maps $H_t := \mathbb{H} \setminus K_t$ conformally onto $\mathbb{H}$ and the inverse mapping $f_t := g_t^{-1}$ satisfies the partial differential equation

$$\partial_t f_t = -f'_t \frac{a}{z - U_t}, \quad f_0(z) = z. \quad (2)$$

Throughout the paper we will use the notation

$$\hat{f}_t(z) := f_t(U_t + z),$$

for $z \in \mathbb{H}$.

The time-reversed Loewner equation

$$\partial_t F_t = -\frac{a}{F_t - U_{T-t}}, \quad t \in (0, T], \quad F_0(z) = z \quad (3)$$

is often useful to avoid dealing directly with (2): it is easy to see that if F is a solution (3) and f a solution to (2) then

$$F_T(z) = f_T(z).$$

If there is a curve $\gamma(t)$ such that H_t is the unbounded connected component of $\mathbb{H} \setminus \gamma[0, t]$ we say that the Loewner chain (g_t) is generated by a curve. This property is known [RS05, Theorem 4.1] to be equivalent to the existence of the radial limit

$$\lim_{y \rightarrow 0} \hat{f}_t(iy) =: \gamma(t) \quad (4)$$

for each $t > 0$ together with continuity of $t \mapsto \gamma(t)$. Loewner chains corresponding to driving functions with strictly higher regularity than Hölder-1/2 are always generated by a simple curve; in fact Hölder-1/2 continuity with a sufficiently small norm is sufficient to guarantee a simple curve. Conversely, there are examples of Loewner chains corresponding to Hölder-1/2 functions (with large norm) that are not generated by a curve, see [MR05] for these results.

In particular we will be interested in Schramm-Loewner evolution, $\text{SLE}(\kappa)$, defined as the Loewner chain corresponding to $a = 2$ and $U_t = \sqrt{\kappa}B_t$, where B is standard Brownian motion and $\kappa \geq 0$. SLE is known to be generated by a curve. We note that the $\text{SLE}(\kappa)$ path is simple if and only if $0 \leq \kappa \leq 4$ and space filling for $\kappa \geq 8$, see [RS05] for proofs of these facts. It may be noted that there is presently no known direct proof of that $\text{SLE}(8)$ is generated by a curve, see [LSW04] for an indirect proof.

Definition 2.1. A *(positive) subpower function* is a continuous, non-decreasing function $\varphi : [0, \infty) \rightarrow (0, \infty)$ that satisfies

$$\lim_{x \rightarrow \infty} x^{-\nu} \varphi(x) = 0$$

for all $\nu > 0$.

3. DETERMINISTIC RESULTS

In this section we prove a number of results about Loewner chains and associated curves that are not special to SLE. In particular we consider Loewner chains corresponding to driving functions which are weakly Hölder-1/2, see (13).

For convenience we state the following well-known result, see [Pom92] for a proof.

Lemma 3.1 (The Koebe distortion and one-quarter theorems). *Suppose $f : D \rightarrow \mathbb{C}$ is a conformal map and set $d = \text{dist}(z, \partial D)$ for $z \in D$. Then*

$$\frac{1-r}{(1+r)^3} |f'(z)| \leq |f'(w)| \leq \frac{1+r}{(1-r)^3} |f'(z)|, \quad |z-w| \leq rd, \quad (5)$$

and

$$B(f(z), d|f'(z)|/4) \subset f(D), \quad (6)$$

where $B(w, \rho)$ denotes the open disk of radius ρ around w .

Lemma 3.2. *Let S be the rectangle $S = \{x + iy : -1 < x < 1, 0 < y < 1\}$. There exist $c, \alpha < \infty$ such that if f is a conformal map defined on $2S$, $z, w \in S$ and $\text{Im } z, \text{Im } w \geq 1/r$, then*

$$|f'(z)| \leq cr^\alpha |f'(w)|. \quad (7)$$

Proof. One way to prove this is to take a conformal transformation of $2S$ onto the unit disk with $f(z) = 0$ and using the distortion theorem. A more direct approach is as follows. We may assume that $\text{Im } z = 1/r$. Take a Whitney decomposition of S , that is, a partition of S into dyadic rectangles $\{S_{j,k}\}$ where

$$S_{j,k} = \{x + iy : j2^{-k} \leq x \leq (j+1)2^{-k}, 2^{-(k+1)} \leq y \leq 2^{-k}\},$$

for $j = 0, \dots, 2^k - 1$, and $k \in \mathbb{N}$. Let u, v be in the same rectangle. Then by iterating (5) it follows that there exists a constant c_1 (uniform for all rectangles; 12^5 would work for instance) such that

$$c_1^{-1} \leq |f'(u)|/|f'(v)| \leq c_1. \quad (8)$$

Suppose that $z \in S_{j,k}$. Then $k \leq \log r \leq k+1$. It follows that there exists a path in S (the hyperbolic geodesic, for instance) that connects z to w and intersects at most $2\lceil \log r \rceil + 2$ rectangles. Hence, there is a constant c_2 such that by iterating (8) along the path at most $c_2(\log r + 1)$ times

$$\begin{aligned} |f'(z)| &\leq c_1^{c_2(\log r + 1)} |f'(w)| \\ &= cr^\alpha |f'(w)|, \end{aligned}$$

for $\alpha = c_2 \log c_1$ and $c = c_1^{c_2}$. $\square$

A proof of the next lemma may be found in [Law05]. For a set K , we let $\text{diam}(K)$ denotes the diameter of K and $\text{height}(K) := \sup\{\text{Im } z : z \in K\}$.

Lemma 3.3. *Suppose K_t is the hull obtained by solving (1) with U_t as driving function. Let $R(t) = \sqrt{t} + \sup_{0 \leq s \leq t} \{|U_s - U_0|\}$. Then there exists a constant $c < \infty$ such that*

$$c^{-1}R(t) \leq \text{diam}(K_t) \leq cR(t). \quad (9)$$

Lemma 3.4. *Let K be a hull. There exists a constant $c < \infty$ such that*

$$\text{hcap}(K) \leq c \text{diam}(K) \text{height}(K). \quad (10)$$

Proof. Let $d = \text{diam}(K)$ and $h = \text{height}(K)$. We shall only consider the case when $h < d$ (the other case is easily verified by considering the map $z \mapsto z + d^2/z$). By scaling, translation invariance, and monotonicity

of hcap we may assume that $d = 1$ and that K is contained in the rectangle $R = \{z : |\operatorname{Re} z| \leq 1/2, 0 \leq \operatorname{Im} z \leq h\}$, so that $\text{hcap}(K) \leq \text{hcap}(R)$. It is known that

$$\text{hcap}(R) = \lim_{y \rightarrow \infty} y \mathbb{E}^{iy}(\operatorname{Im} B(T)),$$

where B denotes a complex Brownian motion and T denotes the hitting time of $\partial(\mathbb{H} \setminus R)$, see [Law05]. Hence

$$\text{hcap}(R) \leq \lim_{y \rightarrow \infty} y \omega(iy, \partial R, \mathbb{H} \setminus R) \cdot h,$$

where ω denotes harmonic measure. Note that R can be covered by $O(h^{-1})$ discs of radius $2h$ centered on the real line. The harmonic measure from iy of any such disc is bounded from above by the harmonic measure of the disc centered at the origin. Since

$$\begin{aligned} \omega(iy, \partial(2h\mathbb{D}), \mathbb{H} \setminus (2h\mathbb{D})) &= (4/\pi) \arctan(2h/y) \\ &\leq ch/y \end{aligned}$$

for large y , the lemma follows from the maximum principle. $\square$

Lemma 3.5. *Suppose f_t satisfies (2) and $z = x + iy \in \mathbb{H}$, then for $s \geq 0$*

$$e^{-5as/y^2} |f'_t(z)| \leq |f'_{t+s}(z)| \leq e^{5as/y^2} |f'_t(z)|. \quad (11)$$

In particular, if $s \leq y^2$,

$$e^{-5a} |f'_t(z)| \leq |f'_{t+s}(z)| \leq e^{5a} |f'_t(z)|.$$

Also, if $0 \leq s \leq y^2$,

$$|f_{t+s}(z) - f_t(z)| \leq \frac{y}{5} [e^{5a} - 1] |f'_t(z)|. \quad (12)$$

Proof. Without loss of generality assume that $a = 1$. Differentiating (2) yields

$$\partial_t f'_t(z) = -f''_t(z) \frac{1}{z - V_t} + f'_t(z) \frac{1}{(z - V_t)^2}.$$

Note that $|z - V_t| \geq y$. Applying Bieberbach's theorem (the $n = 2$ case of the Bieberbach conjecture) to the disk of radius y about z , we can see that

$$|f''_t(z)| \leq 4y^{-1} |f'_t(z)|.$$

and hence

$$|\partial_t f'_t(z)| \leq 5y^{-2} |f'_t(z)|,$$

which implies (11). Returning to (2), we see that

$$|\partial_t f_t(z)| \leq |f'_t(z)| \frac{a}{\operatorname{Im}(z)}.$$

Using (11), we see that

$$\begin{aligned} |f_{t+s}(z) - f_t(z)| &\leq \int_0^s |\partial_s f_{t+s}(z)| ds \\ &\leq \frac{a|f'_t(z)|}{y} \int_0^s e^{5as/y^2} ds = \frac{y}{5} (e^{5a} - 1) |f'_t(z)|. \end{aligned}$$

□

We shall consider Loewner chains corresponding to functions that are Hölder continuous of order α for any $\alpha < 1/2$. We say that U is *weakly Hölder-1/2* if there exists a subpower function φ such that $r^{1/2}\varphi(1/r)$ is a modulus of continuity for U , that is,

$$\sup_{|s| \leq r} |U_{t+s} - U_t| \leq r^{1/2} \varphi(1/r). \quad (13)$$

By P. Lévy's theorem the sample paths of Brownian motion are almost surely weakly Hölder-1/2 with subpower function $c\sqrt{\log(r)}$, $c > \sqrt{2}$, see [RY99, Theorem I.2.7]. Therefore all results for Loewner chains corresponding to functions that satisfy (13) hold for SLE(κ) with probability one.

Lemma 3.6. *There exist constants $c, \alpha < \infty$ such that the following holds. Let (g_t) be a Loewner chain corresponding to the continuous function U_t . Let $s \in [0, y^2]$ for $y > 0$. Then*

$$|f'_{t+s}(U_{t+s} + iy)| \leq cM^\alpha |f'_{t+s}(U_t + iy)|, \quad (14)$$

where $M = \max\{|U_{t+s} - U_t|/y, 1\}$. In particular, if U_t satisfies (13), then there exists a subpower function φ such that for all t and all $s \in [0, y^2]$,

$$|f'_{t+s}(U_{t+s} + iy)| \leq \varphi(1/y) |f'_{t+s}(U_t + iy)|.$$

Proof. We first rescale by $|U_{t+s} - U_t|$, and then apply Lemma 3.2 with $r = \max\{|U_{t+s} - U_t|/y, 1\}$ to get the conclusion. □

The last two lemmas immediately imply the following result, which we record as a lemma. This essentially says that for weakly Hölder-1/2 Loewner chains, it is enough to consider the derivative at dyadic times.

Lemma 3.7. *Let (g_t) be the Loewner chain corresponding to U_t satisfying (13). Suppose that there exist constants c and β such that for all $n \geq 1$*

$$|\hat{f}'_{t_k}(i2^{-n})| \leq c2^{n\beta}, \quad (15)$$

where $t_k = k2^{-2n}$, $k = 0, 1, \dots, 2^{2n}$. Then for every $\beta_1 > \beta$, there exists a constant $c_1 < \infty$ such that

$$|\hat{f}'_t(i2^{-n})| \leq c_1 2^{n\beta_1},$$

for $t \in [0, 1]$.

Let (g_t) be the Loewner chain corresponding to a function U_t satisfying (13) that is generated by a curve $\gamma(t)$. We want to estimate the modulus of continuity of γ . The following quantity will be useful

$$v(t, y) := \int_0^y |f'_t(U_t + ir)| dr, \quad y > 0. \quad (16)$$

The geometrical interpretation is of course the length (if it exists) of the image of the segment $[U_t, U_t + iy]$ under f_t . For a given t , the limit

$$\gamma(t) = \lim_{y \rightarrow 0+} f_t(U_t + iy) \quad (17)$$

exists if $v(t, y)$ is finite for some $y > 0$. By integration we have

$$|\gamma(t) - f_t(U_t + iy)| \leq v(t, y).$$

Using the Koebe one-quarter theorem, we can see that

$$v(t, y) \geq y |\hat{f}'_t(iy)|/4. \quad (18)$$

The next result shows that Loewner chains corresponding to weakly Hölder-1/2 functions are generated by a curve if $v(t, y)$ decays polynomially in y . We also get an estimate of the modulus of continuity of the curve.

Proposition 3.8. *Let (g_t) be the Loewner chain corresponding to U_t satisfying (13). Then there exists a subpower function φ such that if $0 \leq t \leq t + s \leq 1$ and $s \in [0, y^2]$*

$$|\gamma(t + s) - \gamma(t)| \leq \varphi(1/y) [v(t + s, y) + v(t, y)]. \quad (19)$$

Remark. We have not assumed existence of the curve in this proposition. If $v(t + s, y), v(t, y) < \infty$, then we know that radial limit (4) exists, so we can write $\gamma(t), \gamma(t + s)$. If one of the radial limits does not exist we can define γ any way that we want since the right hand side of (19) is infinite.

Proof. We start by writing

$$\begin{aligned} |\gamma(t + s) - \gamma(t)| &\leq |\gamma(t + s) - f_{t+s}(U_{t+s} + iy)| + |\gamma(t) - f_t(U_t + iy)| \\ &\quad + |f_{t+s}(U_{t+s} + iy) - f_{t+s}(U_t + iy)| + |f_{t+s}(U_t + iy) - f_t(U_t + iy)|. \end{aligned}$$

We have to estimate the last two terms. Note that

$$|f_{t+s}(U_{t+s} + iy) - f_{t+s}(U_t + iy)| \leq |U_{t+s} - U_t| \max |f'_{t+s}(w)|,$$

where the maximum is over all w on the line segment connecting $U_{t+s} + iy$ and $U_t + iy$. By Lemma 3.6 using assumption (13) and then by (18), we see that

$$\begin{aligned} |f_{t+s}(U_{t+s} + iy) - f_{t+s}(U_t + iy)| &\leq c \varphi(1/y) y |f'_{t+s}(U_{t+s} + iy)| \\ &\leq c \varphi(1/y) v(t+s, y). \end{aligned}$$

By (12) and (18)

$$\begin{aligned} |f_{t+s}(U_t + iy) - f_t(U_t + iy)| &\leq \frac{y}{5} [e^{5a} - 1] |f'_t(U_t + iy)| \\ &\leq c [e^{5a} - 1] v(t, y). \end{aligned}$$

□

Proposition 3.9. *Let (g_t) be the Loewner chain corresponding to U_t satisfying (13). Suppose that (g_t) is generated by a curve $\gamma(t)$. Then for each t , there exist $t_1, t_2 \in [t, t+y^2]$, and constants $c, \delta > 0$ such that*

$$|\gamma(t_1) - \gamma(t_2)| \geq c |\hat{f}'_t(iy)| y \varphi(1/y)^{-\delta}.$$

Proof. Let $\beta := g_t(\gamma[t, t+y^2])$. Then β is the curve obtained by solving (1) with U_{t+r} , $r \in [0, y^2]$, as driving function. Hence $\text{hcap}(\beta) = ay^2$, and by Lemma 3.3 together with the assumption (13) on U ,

$$\rho_d := \text{diam}(\beta) \leq y \varphi(1/y). \quad (20)$$

Next, we combine (20) with Lemma 3.4 to find

$$\rho_h := \text{height}(\beta) \geq c y \varphi(1/y)^{-1}, \quad (21)$$

for some constant c . Let $z \in \beta$ satisfy $\text{Im } z \geq \rho_h$. Note that $|\text{Re } z - U_t| \leq \text{diam}(\beta) \leq \rho_d$. By scaling by ρ_d^{-1} and then using Lemma 3.2 with $r = \varphi(1/y)^2$ we get

$$|f'_t(z)| \geq c \varphi(1/y)^{-2\alpha} |\hat{f}'_t(iy)|, \quad (22)$$

where α is the exponent from Lemma 3.2. Let w be a point on β such that $|w - z| = \rho_h/2$. Since $z, w \in \beta$ there are $t_1, t_2 \in [t, t+y^2]$ such that $\gamma(t_1) = f_t(z)$ and $\gamma(t_2) = f_t(w)$. In view of (6) we have

$$B(f_t(z), |f'_t(z)| \rho_h/16) \subset f_t(B(z, \rho_h/4)), \quad (23)$$

where $B(z, \rho)$ denotes the open ball of radius ρ around z . Hence, using (23) and (22), we conclude that

$$\begin{aligned} |\gamma(t_1) - \gamma(t_2)| &= |f_t(z) - f_t(w)| \\ &\geq |f'_t(z)| \rho_h/16 \\ &\geq c |\hat{f}'_t(iy)| y \varphi(1/y)^{-(2\alpha+1)}, \end{aligned}$$

and this completes the proof. □

Although we do not use it in this paper, we will give some results about existence of the curve for continuous driving functions that are not necessarily weakly Hölder-1/2.

Proposition 3.10. *There exists $c < \infty$ such that the following is true. Suppose $\delta > 0$ and (g_t) is a Loewner chain with driving function U_t . Suppose that $0 \leq s, y < \infty$ and*

$$|U_{t+s} - U_s| \leq \delta. \quad (24)$$

Then

$$|\gamma(t+s) - \gamma(t)| \leq c e^{5a} [v(t, \delta) + v(t+s, \delta)].$$

Proof. We use the triangle inequality on $|\gamma(t+s) - \gamma(t)|$ as in the beginning of the proof of Proposition 3.8 with $y = \delta$. The first two terms are bounded by $v(t+s, \delta)$ and $v(t, \delta)$, respectively, and the fourth term is bounded in the same way. The distortion theorem, (24), and (18) imply

$$\begin{aligned} |f_{t+s}(U_{t+s} + i\delta) - f_{t+s}(U_t + i\delta)| &\leq c\delta |f'_{t+s}(U_{t+s} + i\delta)| \\ &\leq c v(t+s, \delta), \end{aligned}$$

and we get the desired estimate. $\square$

Corollary 3.11. *Suppose (g_t) is a Loewner chain with continuous driving function U_t . Suppose that for each $0 < t_1 < t_2 < \infty$,*

$$\lim_{y \rightarrow 0+} v(t, y) = 0$$

uniformly for $t \in [t_1, t_2]$. Then (g_t) is generated by a curve.

Proof. Since γ is a uniform limit of a sequence of continuous functions on $[t_1, t_2]$, γ is continuous on $[t_1, t_2]$. One can check directly from the Loewner equation that γ is right continuous at 0 and hence γ is continuous. $\square$

Remark. This result also gives an estimate for the modulus of continuity of γ . However, the assumptions are very strong. If we do not assume that U_t is weakly Hölder-1/2, we do not have Lemma 3.7.

4. PROOF OF THEOREM 1.1

We now turn to the proof of our main result. The proof of Theorem 1.1 is split into two parts: the lower bound (which requires derivative estimates from above) is proven in Subsection 4.1 and the upper bound (which requires estimates from below and control of correlations) in Subsection 4.2. The SLE moment estimates (Lemma 4.1 and Lemma 4.4)

we need for the proof are only stated in this section. The proofs of the estimates build on those in [Law09] and are discussed in the last section.

To state the lemmas it is convenient to introduce a number of κ -dependent parameters. Suppose

$$-\infty < r < r_c := \frac{1}{2} + \frac{4}{\kappa}.$$

The significance of r_c is discussed in Section 5. Let

$$\lambda = \lambda(r) = r \left(1 + \frac{\kappa}{4}\right) - \frac{\kappa r^2}{8},$$

$$\zeta = \zeta(r) = r - \frac{\kappa r^2}{8},$$

and

$$\beta = \beta(r) = -1 + \frac{\kappa}{4 + \kappa - \kappa r}.$$

Note that β and λ strictly increase with r for $-\infty < r < r_c$ and hence, we could alternatively consider either of them as the free parameter. Let r_+ be the larger root to $\lambda\beta + \zeta = 2$ and let β_+ , λ_+ and ζ_+ be the corresponding values of β , λ and ζ respectively. Note that if $\kappa \neq 8$, then $r_+ < r_c$ and

$$-1 = \beta(-\infty) < \beta_+ = -1 + \frac{\kappa}{12 + \kappa - 4\sqrt{8 + \kappa}} < \beta(r_c) = 1.$$

Also,

$$\alpha_* = \frac{1 - \beta_+}{2}.$$

4.1. Lower bound. The following is the main moment estimate for the lower bound. This was proved in [Law09] for a certain range of r including $r = 1, \kappa < 8$ which was most important for that paper. We give a different proof here that is valid for all $r < r_c$.

Lemma 4.1. *Suppose $r < r_c$. Then there exists $c < \infty$ such that for all $t \geq 1$*

$$\mathbb{E} \left[|\hat{f}'_{t^2}(i)|^\lambda \right] \leq c t^{-\zeta}. \quad (25)$$

Proof. See Section 5. □

In the proof of this result one also finds that the expectation in (25), roughly speaking, is carried on an event on which $|\hat{f}'_{t^2}(i)| \approx t^\beta$ and this has probability of order $t^{-(\zeta + \lambda\beta)}$. From this lemma we can derive the following uniform estimate from which the lower bound will follow.

Recall that $\hat{f}'_0(iy) = 1$ for all $y > 0$, so we have to restrict our attention to positive β .

Proposition 4.2. *Suppose $\beta > \max\{0, \beta_+\}$. With probability one there exists $y_0 > 0$, such that for all $t \in [0, 1]$ and all $y < y_0$,*

$$|\hat{f}'_t(iy)| \leq y^{-\beta}.$$

Proof. For $\beta > \max\{0, \beta_+\}$ we have $\lambda > 0, \zeta > 0$ and $\beta\lambda + \zeta > 2$. By choosing β smaller if necessary, but still larger than $\max\{0, \beta_+\}$, we can guarantee that $\zeta < 2$. We write

$$\hat{f}_{j,n} = \hat{f}_{(j-1)2^{-2n}}, \quad j = 1, \dots, 2^{2n}.$$

By Lemma 3.7 and the distortion theorem it suffices to prove that for all $\beta > \max\{0, \beta_+\}$, with probability one there exists $N < \infty$ such that for $n \geq N$,

$$|\hat{f}'_{j,n}(i2^{-n})| \leq 2^{\beta n}, \quad j = 1, 2, \dots, 2^{2n}. \quad (26)$$

Note that scale invariance implies

$$\mathbb{P} \left\{ |\hat{f}'_{j,n}(i2^{-n})| \geq 2^{\beta n} \right\} = \mathbb{P} \left\{ |\hat{f}'_{j-1}(i)| \geq 2^{\beta n} \right\}.$$

By Lemma 4.1 and Chebyshev's inequality,

$$\begin{aligned} \mathbb{P} \left\{ |\hat{f}'_{j-1}(i)| \geq 2^{\beta n} \right\} &\leq 2^{-\lambda\beta n} \mathbb{E} \left[|\hat{f}'_{j-1}(i)|^\lambda \right] \\ &\leq c j^{-\frac{\zeta}{2}} 2^{-\lambda\beta n} = c \left(\frac{j}{2^{2n}} \right)^{-\frac{\zeta}{2}} 2^{-n(\lambda\beta + \zeta)}. \end{aligned}$$

Since $\lambda\beta + \zeta > 2$ and $\zeta < 2$, we can sum over j to get

$$\sum_{j=1}^{2^{2n}} \mathbb{P} \left\{ |\hat{f}'_{j-1}(i)| \geq 2^{\beta n} \right\} \leq c 2^{-n(\lambda\beta + \zeta - 2)}, \quad (27)$$

and hence

$$\sum_{n=1}^{\infty} \sum_{j=1}^{2^{2n}} \mathbb{P} \left\{ |\hat{f}'_{j-1}(i)| \geq 2^{\beta n} \right\} < \infty.$$

The Borel-Cantelli lemma now implies (26). $\square$

Proposition 4.3. *Suppose $\beta_+ < \beta < 0$. With probability one, for every $\epsilon > 0$, there exists $y_\epsilon > 0$ such that for all $t \in [\epsilon, 1]$ and all $y < y_\epsilon$,*

$$|f'_t(iy)| \leq y^{-\beta}.$$

Proof. For $\beta_+ < \beta < 0$, we have $\lambda\beta + \zeta > 2$ and $\lambda > 0$. The proof is identical to the previous proposition except that we do not have $\zeta > 0$. We replace (27) with

$$\sum_{\epsilon 2^{2n} \leq j \leq 2^{2n}} \mathbb{P} \left\{ |\hat{f}'_{j-1}(i)| \geq 2^{\beta n} \right\} \leq c_\epsilon 2^{-n(\lambda\beta + \zeta - 2)}.$$

□

We can now easily prove the lower bound in Theorem 1.1. Recall that $\alpha_* = \min\{1/2, (1 - \beta_+)/2\}$.

Proof of lower bound for Theorem 1.1. Let $1 \geq \beta > \tilde{\beta} > \max\{0, \beta_+\}$; recall that $\beta_+ = 1$ if and only if $\kappa = 8$. Almost surely, by Proposition 4.2, we have for all $t \in [0, 1]$

$$v(t, y) = \int_0^y |\hat{f}'_t(ir)| dr \leq y^{1-\tilde{\beta}} \quad (28)$$

if y is small enough. The last inequality together with Proposition 3.8 show that the SLE(κ) Loewner chain is generated by a curve when $\kappa \neq 8$ and imply the following modulus of continuity

$$|\gamma(t+s) - \gamma(t)| \leq cs^{(1-\beta)/2}$$

for all s small enough since $\tilde{\beta} > \beta$. The lower bound follows.

If $\beta_+ < \beta < 0$, it suffices to prove the result for each fixed ϵ . The argument is the same using Proposition 4.3 □

4.2. Upper bound. In this subsection we shall use the notation

$$\hat{f}_{j,n} = \hat{f}_{(j-1)/n^2}, \quad j = n^2/2, \dots, n^2.$$

Lemma 4.4. *Suppose $r < r_c$. Then there exist $0 < c_1, c_2 < \infty$, a subpower function φ , and events*

$$E_{j,n}, \quad n = 1, 2, \dots, \quad j = 1, \dots, n^2$$

such that the following hold. Let $E(j, n) = 1_{E_{j,n}}$ and

$$F(j, n) = n^{\zeta-2} |\hat{f}'_{j,n}(i/n)|^\lambda E(j, n).$$

- *If $n^2/2 \leq j \leq n^2$, then on the event $E_{j,n}$,*

$$|\hat{f}'_{j,n}(i/n)| \geq \varphi(n)^{-1} n^\beta. \quad (29)$$

- *If $n^2/2 \leq j \leq n$,*

$$c_1 n^{-2} \leq \mathbb{E}[F(j, n)] \leq c_2 n^{-2}. \quad (30)$$

- If $n^2/2 \leq j \leq k \leq n^2$,

$$\mathbb{E}[F(j, n) F(k, n)] \leq n^{-4} \left(\frac{n^2}{k-j+1} \right)^{\frac{\lambda\beta+\zeta}{2}} \varphi \left(\frac{n^2}{k-j+1} \right). \quad (31)$$

Proof. See Section 5.2. $\square$

For fixed β , we let $A_n = A_{n,\beta}$ denote the event that there exists an integer j with $1 \leq j \leq n^2$,

$$|\hat{f}'_{j,n}(i/n)| \geq \varphi(n)^{-1} n^\beta, \quad (32)$$

where φ is as in (29). We then have the following.

Lemma 4.5. *Suppose $\lambda\beta + \zeta < 2$. Then there exist $c > 0$ such for all n sufficiently large,*

$$\mathbb{P}(A_n) \geq c.$$

In particular,

$$\mathbb{P}\{A_n \text{ i.o.}\} \geq c.$$

Proof. Let $F(j, n)$ be as in Lemma 4.4, and let

$$Y_n = \sum_{j=n^2/2}^{n^2} F(j, n).$$

Note that $A_n \supset \{Y_n > 0\}$. The estimates from Lemma 4.4 show that there exist $0 < c < c_2 < \infty$ such that

$$\mathbb{E}[Y_n] \geq c, \quad \mathbb{E}[Y_n^2] \leq c_2^2.$$

(This uses $\beta\lambda + \zeta < 2$.) Therefore, a standard second moment argument gives

$$\mathbb{P}(A_n) \geq \mathbb{P}\{Y_n > 0\} \geq \frac{\mathbb{E}[Y_n]^2}{\mathbb{E}[Y_n^2]} \geq \frac{c^2}{c_2^2}.$$

$\square$

We can now prove the upper bound for Theorem 1.1 and thereby complete the proof. Notice that Proposition 3.9 immediately implies that $\gamma[0, t]$ cannot be Hölder continuous of order $> 1/2$, since $\hat{f}'_0(z) = 1$.

Proof of upper bound for Theorem 1.1. Let $\beta < \tilde{\beta} < \beta_+$. Proposition 3.9 implies that on the event $A_n = A_{n,\tilde{\beta}}$ there exist times $t_1, t_2 \in [0, 1]$ such that

$$|\gamma(t_1) - \gamma(t_2)| \geq cn^{\beta-1} = c(n^{-2})^{(1-\beta)/2}, \quad |t_1 - t_2| \leq n^{-2}.$$

Consequently on the event $\{A_n \text{ i.o.}\}$, the curve $\gamma(t)$, $t \in [0, 1]$, is not Hölder continuous of order $(1-\beta)/2$. To show that this happens almost surely, let $\mathcal{A}_r$ be the event that $\gamma(t)$, $t \in [0, r]$, is not Hölder continuous of order $(1-\beta)/2$. By Lemma 4.5 we have

$$\mathbb{P}(\mathcal{A}_1) =: c_0 > 0,$$

and by scale invariance

$$\mathbb{P}(\mathcal{A}_r) = c_0, \quad r > 0.$$

Note that $\mathcal{A}_r \subset \mathcal{A}_1$ if $r \leq 1$ so that

$$\mathbb{P}\left(\bigcap_{r>0} \mathcal{A}_r\right) = c_0.$$

Since $c_0 > 0$, it now follows from the Blumenthal zero-one law (see [RY99, Theorem III.2.15]) that $c_0 = 1$ and this completes the proof. $\square$

5. MOMENTS OF DERIVATIVES

In this section we review some results from [Law09] and extend one result. We fix κ and we let $a = 2/\kappa$. We also fix a real number r such that

$$r < r_c = \frac{1+4a}{2}.$$

All constants and parameters in this section depend on a and r . We let

$$q = r_c - r = 2a + \frac{1}{2} - r > 0.$$

The positivity of q is important for the arguments in this section and this is why r must be less than r_c .

A useful tool for estimating moments of $|\hat{f}'_t|$ is the reverse Loewner flow (see, e.g., [Law09, Section 10.3]). Suppose U_t is a standard Brownian motion and $h_t(z)$ is the solution to the reverse-time Loewner equation

$$\partial_t h_t(z) = \frac{a}{U_t - h_t(z)}, \quad h_0(z) = z. \quad (33)$$

For fixed T , the distribution of $h_T(z) - U_T$ is the same as that of $\hat{f}_T(z)$ and hence, $h'_T(z)$ has the same distribution as $\hat{f}'_T(z)$. Indeed, suppose g_t is the solution of the forward-time equation for some continuous V with $V_0 = 0$:

$$\partial_t g_t(z) = \frac{a}{g_t(z) - V_t},$$

and we let $V_t^{(T)} = V_{T-t} - V_T$. Then if h is a solution to (33) with driving function $V_t^{(T)}$ we have

$$g_T^{-1}(z + V_T) = h_T(z) + V_T = h_T(z) - V_T^{(T)}.$$

It remains to note that $V_t^{(T)}$, $0 \leq t \leq T$, is a standard Brownian motion starting at 0 if V is. If $S < T$ and $h^{(S)}, h^{(T)}$ denote the solutions to (33) with driving functions $U_t^{(S)} = U_{S-t} - U_S$ and $U_t^{(T)} = U_{T-t} - U_T$ then $h_t^{(T)}$, $0 \leq t \leq T - S$ and $h_t^{(S)}$, $0 \leq t \leq S$ are independent. Note that $\hat{f}'_S(z) \hat{f}'_T(z)$ has the same distribution as $(h_S^{(S)})'(z) (h_T^{(T)})'(z)$.

Let $Z_t = Z_t(i) = X_t + iY_t = h_t(i) - U_t$ and $S_t = \sin[\arg Z_t] = [1 + X_t^2/Y_t^2]^{-1/2}$. To study the behavior of $h'_t(i)$, it is useful to do a time-change so that the the logarithm of the imaginary part grows linearly. Let

$$\sigma(t) = \inf\{s : Y_s(z) = e^{at}\}.$$

The following lemma can be deduced from the Loewner equation and the analogous time change in a stochastic differential equation.

Lemma 5.1. [Law09, Section 5] *Suppose J_t satisfies*

$$dJ_t = -r_c \tanh J_t dt + dW_t, \quad J_0 = 0, \quad (34)$$

where W_t is standard Brownian motion. Let

$$L_t = t - \int_0^t \frac{2 ds}{\cosh^2 J_s},$$

$$\sigma(t) = \int_0^t e^{2as} \cosh^2 J_s ds, \quad \eta(s) = \sigma^{-1}(s). \quad (35)$$

Then the joint distribution of

$$e^{aL_{\eta(s)}}, \quad e^{a\eta(s)}, \quad \cosh J_{\eta(s)}, \quad 0 \leq s < \infty$$

is the same as that of

$$|h'_s(i)|, \quad Y_s, \quad S_s^{-1}, \quad 0 \leq s < \infty.$$

Let λ, ζ, β be as in Section 4; we can write

$$\lambda = r \left(1 + \frac{1}{2a}\right) - \frac{r^2}{4a}, \quad \zeta = \lambda - \frac{r}{2a},$$

$$\beta = \frac{1 - 2q}{1 + 2q} = \frac{r - 2a}{1 - 2r + 2a}.$$

We comment here that the β of this paper is the same as μ in [Law09]; the β in that paper is half this value.

Note that (34) can be written as

$$dJ_t = -(q + r) \tanh J_t dt + dW_t, \quad J_0 = 0.$$

Using Itô's formula, one can see that

$$N_t = e^{a\lambda L_t} e^{a\zeta t} [\cosh J_t]^r$$

is a martingale satisfying

$$dN_t = r [\tanh J_t] N_t dW_t.$$

Let $\mathbb{P}_*$ denote the probability measure obtained by weighting by the martingale N_t , that is, if E is an event measurable with respect to $\{W_s : 0 \leq s \leq t\}$,

$$\mathbb{P}_*(E) = N_0^{-1} \mathbb{E}[N_t 1_E] = \mathbb{E}[N_t 1_E].$$

The Girsanov theorem (see, e.g., [RY99, Chapter VIII] and [Law09, Appendix A]) implies that

$$\begin{aligned} dW_t &= r \tanh J_t dt + dB_t, \\ dJ_t &= -q \tanh J_t dt + dB_t, \end{aligned} \tag{36}$$

where B_t is a standard Brownian motion with respect to the measure $\mathbb{P}_*$. We write $\mathbb{E}_*$ for expectations with respect to $\mathbb{P}_*$.

Lemma 5.2. [Law09, Lemma 7.1] *Suppose J_t satisfies (36).*

- J_t is a positive recurrent diffusion (with respect to the measure $\mathbb{P}_*$) with invariant density

$$v(x) = \frac{\Gamma(q + \frac{1}{2})}{\Gamma(\frac{1}{2})\Gamma(q)} \frac{1}{\cosh^{2q} x}.$$

•

$$\int_{-\infty}^{\infty} \left[1 - \frac{2}{\cosh^2 x} \right] v(x) dx = \frac{1 - 2q}{1 + 2q} = \beta.$$

- Assume $J_0 = 0$. There exists $c < \infty$ such that if $k \geq 0$, $u \geq 0$,

$$\mathbb{P}_*\{\cosh J_t \geq u \text{ for some } k \leq t \leq k + 1\} \leq c u^{-2q}. \tag{37}$$

Lemma 5.3. *There exists $c < \infty$ such that the following holds. Suppose J_t satisfies (36) with $J_0 = x > 0$.*

- If $y > x$,

$$\mathbb{P}_* \left\{ \max_{0 \leq s \leq 1} |J_s| \geq y \right\} \leq c \exp \left\{ -\frac{(y - x)^2}{2} \right\}. \tag{38}$$

- If $y < x$ and $1 \leq t < (x - y)/q$,

$$\mathbb{P}_* \left\{ \min_{0 \leq s \leq t} J_s \leq y \right\} \leq c \exp \left\{ -\frac{(x - qt - y)^2}{2t} \right\}. \tag{39}$$

Proof. Since the drift of J_t points towards the origin, the distribution of $|J_t|$ is stochastically dominated by the absolute value of a Brownian motion. Hence by the reflection principle (see, e.g., [RY99, Proposition III.3.7]) if V is a standard Brownian motion

$$\begin{aligned} \mathbb{P}_* \left\{ \max_{0 \leq s \leq 1} |J_s| \geq y \right\} &\leq \mathbb{P}_* \left\{ \max_{0 \leq s \leq 1} |V_s| \geq y - x \right\} \\ &\leq 4 \mathbb{P}_* \{V_1 \geq y - x\}, \end{aligned}$$

which gives the first estimate. Also, $J_t \geq \tilde{J}_t$ where $\tilde{J}_t$ satisfies

$$d\tilde{J}_t = -q dt + dB_t.$$

Note that $W_t := \tilde{J}_t + tq - x$ is a standard Brownian motion starting at the origin. We get for $y < x$ and $1 \leq t < (x - y)/q$, again using the reflection principle,

$$\begin{aligned} \mathbb{P}_* \left\{ \min_{0 \leq s \leq t} J_s \leq y \right\} &\leq \mathbb{P}_* \left\{ \max_{0 \leq s \leq t} W_s \geq x - y - qt \right\} \\ &= 2 \mathbb{P}_* \{W_t \geq x - y - qt\}, \end{aligned}$$

and the second estimate follows. $\square$

5.1. Proof of Lemma 4.1. Lemma 4.1 extends a result in [Law09] where the result was proved for a certain range of r (including $r = 1, a > 1/4$ which was most important for that paper). We start by restating it in terms of h with an added lower bound (and upgrading it to a theorem).

Theorem 5.4. *If $r < r_c$, there exists $0 < c_1, c_2 < \infty$ such that for all $t \geq 1$,*

$$c_1 t^{-\zeta} \leq \mathbb{E} [|h'_{t^2}(i)|^\lambda] \leq c_2 t^{-\zeta}.$$

The heuristic argument is fairly straightforward, so let us consider that first. Consider the martingale N_t which we can write as

$$N_t = |h'_{\sigma(t)}(i)|^\lambda e^{at\zeta} S_{\sigma(t)}^{-r}.$$

We know that $\mathbb{E}[N_t] = 1$. If $r < r_c$, then under $\mathbb{P}_*$, since J is positive recurrent, $S_{\sigma(s)}$ tends to be of order 1 for $s < t$. Hence we would expect that $\sigma(t) \approx e^{2at}$ and hence we would expect

$$\mathbb{E} [|h'_{e^{2at}}(i)|^\lambda e^{at\zeta}] \asymp 1.$$

We will only prove the upper bound in Theorem 5.4 in this subsection; the lower bound follows from the work of the next subsection, which uses the upper bound. The next lemma gives a quantitative bound on

the assertion $\sigma(t) \approx e^{2at}$. As a slight abuse of notation, in this section, if E is an event we also write E for the indicator function of the event.

Lemma 5.5. *There exists $c < \infty$ such that for all $u > 0$,*

$$\mathbb{P}_* \{ \sigma(t) \geq u^2 e^{2at} \} \leq c u^{-2q}. \quad (40)$$

Proof. Suppose $\cosh J_s \leq u e^{a(t-s)} (t-s+1)^{-1}$ for all $0 \leq s \leq t$. Then,

$$\sigma(t) = \int_0^t e^{2as} \cosh^2 J_s ds \leq u^2 e^{2at} \int_0^t (t-s+1)^{-2} ds < u^2 e^{2at}.$$

Therefore,

$$\mathbb{P}_* \{ \sigma(t) \geq u^2 e^{2at} \} \leq \mathbb{P}_* \left[\bigcup_{k=1}^{\infty} K_k \right] \leq \sum_{k=1}^{\infty} \mathbb{P}_*(K_k),$$

where $K_k = K_{k,t}$ denotes the event

$$K_k = \{ \cosh J_{t-s} \geq u e^{a(k-1)} (k+1)^{-1} \text{ for some } k-1 \leq s < k \}. \quad (41)$$

By (37),

$$\mathbb{P}_*(K_k) \leq c u^{-2q} e^{-2aqk} k^{2q}, \quad (42)$$

and hence we can sum over k to get the result. $\square$

The next lemma is a useful “smoothing” result that lets us consider the average of $\mathbb{E}[|h'_s(i)|^\lambda]$ over $t^2 \leq s \leq 2t^2$ instead of $\mathbb{E}[h'_{t^2}(i)|^\lambda]$.

Lemma 5.6. *There exists $c < \infty$ such that*

$$\begin{aligned} \mathbb{E} [|h'_{t^2}(i)|^\lambda] &\leq \frac{c}{t^2} \int_0^\infty e^{2as} \mathbb{E} [(\cosh^2 J_s) |h'_{\sigma(s)}(i)|^\lambda I_s] ds \\ &= \frac{c}{t^2} \int_0^\infty e^{as(2-\zeta)} \mathbb{E}_* [(\cosh J_s)^{2-r} I_s] ds, \end{aligned}$$

where $I_s = I_{s,t}$ denotes the event

$$I_s = \{t^2 \leq \sigma(s) \leq 2t^2\}.$$

Proof. By scaling and the distortion theorem, if $t^2 \leq u \leq 2t^2$,

$$\mathbb{E} [|h'_u(i)|^\lambda] = \mathbb{E} [|h'_{t^2}(it/\sqrt{u})|^\lambda] \geq c \mathbb{E} [|h'_{t^2}(i)|^\lambda].$$

Therefore,

$$\mathbb{E} [|h'_{t^2}(i)|^\lambda] \leq \frac{c}{t^2} \int_{t^2}^{2t^2} \mathbb{E} [|h'_u(i)|^\lambda] du.$$

If we let $s = \sigma^{-1}(u)$, we can change variables and write

$$\int_{t^2}^{2t^2} |h'_u(i)|^\lambda du = \int_0^\infty |h'_{\sigma(s)}(i)|^\lambda I_s du = \int_0^\infty e^{2as} (\cosh^2 J_s) |h'_{\sigma(s)}(i)|^\lambda I_s ds.$$

By taking expectations we get the inequality in the lemma, and the equality follows from the definition of $\mathbb{E}_*$. $\square$

Since

$$\sigma(t) \geq \int_0^t e^{2as} ds = \frac{1}{2a} [e^{2at} - 1],$$

there is a c such that $I_{s, e^{at}}$ is empty if $s \geq t + c$. Hence the previous proposition implies that there exists a $c < \infty$ such that

$$\begin{aligned} & \mathbb{E} [|h_{e^{2at}}'(i)|^\lambda] \leq \\ & c e^{-2at} \int_0^{t+c} e^{as(2-\zeta)} \mathbb{E}_* [(\cosh J_s)^{2-r}; e^{2at} \leq \sigma(s) \leq 2e^{2at}] ds. \end{aligned} \quad (43)$$

Lemma 5.7. *There exists $c < \infty$ such that if $u \geq 1$,*

$$\mathbb{E}_*[(\cosh J_t)^{2-r}; u^2 \leq e^{-2at} \sigma(t) \leq 2u^2] \leq c u^{(2-r)+-2q}.$$

Remark. Roughly speaking, we expect that if $\sigma(t) \asymp u^2 e^{2at}$, then $\cosh J_t \approx u$. Hence, we would guess

$$\begin{aligned} & \mathbb{E}_*[(\cosh J_t)^{2-r}; u^2 \leq e^{-2at} \sigma(t) \leq 2u^2] \approx \\ & u^{2-r} \mathbb{P}_* \{u^2 \leq e^{-2at} \sigma(t) \leq 2u^2\} \leq c u^{2-r-2q}, \end{aligned}$$

where the probability is estimated by (40). This lemma makes the argument rigorous but only gives a weaker result for $r > 2$. Lemma 5.8 below establishes the stronger result for some values of $r > 2$.

Proof. Let $I_t = I_{t,u}$ be the event $\{u^2 \leq e^{-2at} \sigma(t) \leq 2u^2\}$. We claim that I_t is contained in the event

$$A = A_{t,u} = \left\{ \min_{(t-1) \vee 0 \leq s \leq t} \cosh^2 J_s \leq 2u^2 e^{2a} \right\}$$

Indeed, this is obvious for $t \leq 1$ since $\cosh J_0 = 1$ and if $t > 1$ and $\cosh^2 J_s > 2e^{2a}u^2$ for $t-1 \leq s \leq t$, then

$$\sigma(t) \geq \int_{t-1}^t e^{2as} \cosh^2 J_s ds > 2u^2 e^{2at}.$$

Let

$$V_k = V_{k,t,u} = I_t \cap \{u e^{(k-1)} \leq \cosh J_t < u e^k\}.$$

Then,

$$\mathbb{E}_*[(\cosh J_t)^{2-r} I_t] = \sum_{k=-\infty}^{\infty} \mathbb{E}_*[(\cosh J_t)^{2-r} V_k].$$

Note that

$$\mathbb{E}_*[(\cosh J_t)^{2-r} V_k] \asymp u^{2-r} e^{k(2-r)} \mathbb{P}_*(V_k). \quad (44)$$

We will first show that

$$\sum_{k=1}^{\infty} \mathbb{E}_*[(\cosh J_t)^{2-r} V_k] \leq c u^{2-r-2q}. \quad (45)$$

Let k_0 be an integer such that $e^{k-1} > 2e^{2a}$ for $k \geq k_0$. Then,

$$\begin{aligned} \sum_{k=1}^{k_0} \mathbb{E}_*[(\cosh J_s)^{2-r} V_k] &\asymp u^{2-r} \sum_{k=1}^{k_0} \mathbb{P}_*(V_k) \\ &\leq u^{2-r} \mathbb{P}_*\{\cosh^2 J_t \geq u^2\} \leq c u^{2-r-2q}. \end{aligned}$$

The last inequality uses (37).

If $k > k_0$, let

$$\eta_t = \inf \{s \geq (t-1) \vee 0 : \cosh^2 J_s = 2e^{2a}u^2\}.$$

Since $V_k \subset A$ and $\cosh^2 J_t > 2e^{2a}u^2$ on V_k , we know that on the event V_k , $t-1 \leq \eta_t < t$. Hence we can estimate

$$\mathbb{P}_*(V_k) \leq \mathbb{P}_*\{\eta_t < t\} \mathbb{P}_*\{\cosh^2 J_t \geq u^2 e^{k-1} \mid \eta_t < t\}.$$

By (37),

$$\mathbb{P}_*\{\eta_t < t\} \leq \mathbb{P}_*\{\cosh^2 J_s \geq 2e^{2a}u^2 \text{ for some } t-1 \leq s \leq t\} \leq c u^{-2q}.$$

Using (38), we can see that there exist c, α such that

$$\mathbb{P}_*\{\cosh^2 J_t \geq u^2 e^{k-1} \mid \eta_t < t\} \leq c e^{-\alpha k^2}.$$

Hence, plugging into (44), we have

$$\mathbb{E}_*[(\cosh J_t)^{2-r} V_k] \leq c u^{2-r-2q} e^{k(2-r)} e^{-\alpha k^2}.$$

Therefore,

$$u^{2q+r-2} \sum_{k=1}^{\infty} \mathbb{E}_*[(\cosh J_t)^{2-r} V_k] \leq c \sum_{k=0}^{\infty} e^{k(2-r)} e^{-\alpha k^2} < \infty.$$

This proves (45).

Let $\tilde{V} = \bigcup_{k=0}^{\infty} V_{-k}$. On the event $\tilde{V}$,

$$1 \leq \cosh J_t \leq u.$$

Hence $(\cosh J_t)^{2-r} \leq u^{(2-r)_+}$ and, using (40),

$$\mathbb{E}_*[(\cosh J_t)^{2-r} \tilde{V}] \leq u^{(2-r)_+} \mathbb{P}_*(\tilde{V}) \leq u^{(2-r)_+} \mathbb{P}_*(I_t) \leq c u^{(2-r)_+} u^{-2q}.$$

□

Lemma 5.8. *If $r > a + 1$, there exists $c < \infty$ such that for all u ,*

$$\mathbb{E}_*[(\cosh J_t)^{2-r}; u^2 \leq e^{-2at} \sigma(t) \leq 2u^2] \leq c u^{2-r-2q}.$$

Proof. We use the notation from the proof of Lemma 5.7 and let $V^j = V_{-j}$. We note that (45) holds for all values of r . Hence, we only need to show

$$\sum_{j=0}^{\infty} \mathbb{E}_*[(\cosh J_t)^{2-r} V^j] \leq c u^{2-r-2q}.$$

Let K_k be as in (41). Then

$$\sum_{j=0}^{\infty} \mathbb{E}_*[(\cosh J_t)^{2-r} V^j] \leq \sum_{k=1}^{\infty} \sum_{j=0}^{\infty} \mathbb{E}_*[(\cosh J_t)^{2-r} V^j K_k].$$

Note that

$$\begin{aligned} u^{2q+r-2} \mathbb{E}_*[(\cosh J_t)^{2-r} V^j K_k] &\asymp u^{2q} e^{j(r-2)} \mathbb{P}_*(V^j \cap K_k) \\ &= u^{2q} \mathbb{P}_*(K_k) e^{j(r-2)} \mathbb{P}_*(V^j \mid K_k). \end{aligned}$$

By (42), we know that $u^{2q} \mathbb{P}_*(K_k) \leq c e^{-2aqk} k^{2q}$. Hence, it suffices to prove that

$$\sum_{k=1}^{\infty} e^{-2aqk} k^{2q} \sum_{j=0}^{\infty} e^{j(r-2)} \mathbb{P}_*(V^j \mid K_k) < \infty. \quad (46)$$

Note that for $1 + a < r < r_c$, we have $a > q$ and

$$-2aq + \frac{(r-2)^2}{2} + (2-r)(a-q) < 0,$$

Choose $\hat{a}$ satisfying $q < \hat{a} < a$ and

$$-2\hat{a}q + \frac{(r-2)^2}{2} + (2-r)(\hat{a}-q) < 0, \quad (47)$$

We will show

$$\sum_{k=1}^{\infty} e^{-2\hat{a}qk} \sum_{j=0}^{\infty} e^{j(r-2)} \mathbb{P}_*(V^j \mid K_k) < \infty.$$

On the event K_k , let s be the largest number less than or equal to k such that

$$\cosh J_{t-s} \geq u e^{a(k-1)} (k+1)^{-1}.$$

By the definition of K_k , we know that $k-1 \leq s \leq k$. Also, we can find c_1 such that

$$J_{t-s} \geq \log u + a(k-1) - \log(k+1) \geq \log u + \hat{a}k - c_1 + \log 2.$$

On the event V^j , $\cosh J_t \leq e^{-j}u$ which implies

$$J_t \leq \log u - j + \log 2.$$

We estimate $\mathbb{P}_*(V^j \mid K_k)$ from above by

$$\sup_{k-1 \leq s \leq k} \mathbb{P}_* \{J_t \leq \log u - j + \log 2 \mid J_{t-s} = \log u + \hat{a}k - c_1 + \log 2\}.$$

As in (39), this probability is bounded above by the corresponding probability if J_t is a Brownian motion with drift $-q$. In particular, (39) shows that there exists M and c_2 such that for $j > M$,

$$\mathbb{P}_*(V^j \mid K_k) \leq c \exp \left\{ -\frac{1}{2k} [(\hat{a} - q)k + j - c_1]^2 \right\}. \quad (48)$$

Note that

$$\sum_{k=1}^{\infty} e^{-2\hat{a}qk} \sum_{j=0}^M e^{j(r-2)} \mathbb{P}_*(V^j \mid K_k) \leq \sum_{k=1}^{\infty} e^{-2\hat{a}qk} \sum_{j=0}^M e^{j(r-2)} < \infty.$$

Also, (48) gives

$$\sum_{j=M+1}^{\infty} e^{j(r-2)} \mathbb{P}_*(V^j \mid K_k) \leq c \sum_{j=M+1}^{\infty} \exp \left\{ j(r-2) - \frac{1}{2k} [j + b]^2 \right\}.$$

where

$$b = (\hat{a} - q)k - c_2.$$

We bound the right hand side from above by a constant times

$$\begin{aligned} \int_{-\infty}^{\infty} e^{x(r-2)} e^{-\frac{(x+b)^2}{2k}} dx &= e^{b(2-r)} \int_{-\infty}^{\infty} e^{y(r-2)} e^{-\frac{y^2}{2k}} dy \\ &\leq c \sqrt{k} \exp \left\{ b(2-r) + \frac{k(r-2)^2}{2} \right\} \\ &\leq c \sqrt{k} \exp \left\{ (2-r)(\hat{a} - q)k + \frac{k(r-2)^2}{2} \right\} \end{aligned}$$

Hence

$$e^{-2\hat{a}qk} \sum_{j=M+1}^{\infty} e^{j(r-2)} \mathbb{P}_*(V^j \mid K_k) \leq C_{\epsilon} \sqrt{k} e^{k\xi}$$

where

$$\xi = -2\hat{a}q + (2-r)(\hat{a} - q) + \frac{(r-2)^2}{2}.$$

Recalling from (47) that $\xi < 0$, we conclude

$$\sum_{k=1}^{\infty} e^{-2\hat{a}qk} \sum_{j=M+1}^{\infty} e^{j(r-2)} \mathbb{P}_*(V^j \mid K_k) < \infty.$$

□

Lemma 5.9. *Suppose $r < r_c$. Then there exists $\theta < 2 - \zeta$ and $c < \infty$ such that for all u*

$$\mathbb{E}_*[(\cosh J_t)^{2-r}; u^2 \leq e^{-2at} \sigma(t) \leq 2u^2] \leq c u^\theta. \quad (49)$$

Proof. We set

$$\theta = \begin{cases} 2 - r - 2q, & \text{if } r \leq 2 \text{ or } r \geq a + \frac{1}{2} \\ -2q & \text{otherwise.} \end{cases}$$

The estimate (49) follows from the previous two lemmas so we only need to verify that $\theta < 2 - \zeta$. It is easy to see that $1 - r - 2q < 2 - \zeta$ for all r . Also, one can check that $-2q < 1 - 4a + r$ provided that $a < 3/5$ or $a \geq 3/5$ and

$$r < 2a \left[3 - \sqrt{5 - (3/a)} \right]. \quad (50)$$

If $a \leq 4$, then one can show that (50) holds for all $r < r_c$. Hence we only need to consider $a \geq 4$ and $r < r_c$ that do not satisfy (50). Such an r satisfies

$$r \geq 2a \left[3 - \sqrt{5 - (3/a)} \right] \geq 2a \left[3 - \sqrt{17/4} \right] \geq 9a/5.$$

In particular, $r > a + 1$. □

Proof of Theorem 5.4. Let θ be as in the previous lemma. Then by (49),

$$\mathbb{E}_*[(\cosh J_s)^{2-r}; e^{2at} \leq \sigma(s) \leq 2e^{2at}] \leq c e^{(t-s)a\theta}.$$

Therefore, by (43),

$$\begin{aligned} \mathbb{E} [|h'_{e^{2at}}(i)|^\lambda] &\leq c e^{-2at} \int_0^{t+c} e^{2as} e^{-as\zeta} e^{(t-s)a\theta} ds \\ &\leq c e^{-a\zeta t} \int_0^{t+c} e^{a(t-s)[-2+\zeta+\theta]} ds. \\ &\leq c e^{-a\zeta t} \int_{-c}^\infty e^{ay[-2+\zeta+\theta]} dy \leq c e^{-a\zeta t}. \end{aligned}$$

The last inequality uses $\theta < 2 - \zeta$. □

5.2. Proof of Lemma 4.4. We essentially follow the proof in [Law09]. In that paper, it was assumed that $r \geq 0$, which we do not want to assume here, but with the upper bound of Theorem 5.4, we can do the argument for all $r < r_c$. We emphasize that the positive recurrence of J_t is critical for this argument and hence we need $q = r_c - r > 0$. Since J_t is positive recurrent, we expect that $J_t = O(1)$ and that (approximately)

$$L_t = \beta t + O(t^{1/2}).$$

The next lemma gives an estimate of this type.

Lemma 5.10. [Law09, Proposition 7.3] *For each $u, t > 0$, let $E_{t,u}$ be the event that the following holds for all $0 \leq s \leq t$:*

$$\begin{aligned} |J_s| &\leq u \log [\min\{s+2, t-s+2\}], \\ |L_s - s\beta| &\leq u \sqrt{s} \log(s+2), \\ |L_t - L_s - (t-s)\beta| &\leq u \sqrt{t-s} \log(t-s+2). \end{aligned}$$

Then,

$$\liminf_{u \rightarrow \infty} \mathbb{P}_*(E_{t,u}) = 1.$$

By integrating, we can see there exists c_0 and $c(u)$ such that for all $t \geq 1$, on the event $E_{t,u}$

$$c_0 e^{2at} \leq \sigma(t) \leq c(u) e^{2at}. \quad (51)$$

Note that the lower bound does not depend on u and follows from the estimate $\cosh^2 J_s \geq 1$. An event such as this is used to define the event $E_t = E_{t,u}$ in Lemma 4.2 where u is sufficiently large so that

$$\mathbb{P}_*(E_t) = \mathbb{E}[N_t \mathbf{1}_{E_t}] \geq \frac{1}{2}.$$

Note that on the event E_t ,

$$|h'_{\sigma(t)}(i)| \asymp e^{at\beta},$$

and using (51), one can show that $|h'_{e^{2at}}(i)| \asymp (e^{at})^\beta$. In fact, there is a function ψ with $\psi(s) = o(s)$ as $s \rightarrow \infty$, such that for all $0 \leq s \leq t$,

$$\begin{aligned} e^{a\beta s - \psi(s)} &\leq |h'_{\sigma(s)}(i)| \leq e^{a\beta s + \psi(s)}, \\ e^{a\beta(t-s) - \psi(t-s)} &\leq \left| \frac{h'_{\sigma(t)}(i)}{h'_{\sigma(s)}(i)} \right| \leq e^{a\beta(t-s) + \psi(t-s)}. \end{aligned}$$

As in (51), we can see that this implies (with perhaps a different $o(s)$ function ψ)

$$\begin{aligned} e^{a\beta s - \psi(s)} &\leq |h'_{e^{2as}}(i)| \leq e^{a\beta s + \psi(s)}, \\ e^{a\beta(t-s) - \psi(t-s)} &|h'_{e^{2as}}(i)| \leq |h'_{e^{2at}}(i)| \leq e^{a\beta(t-s) + \psi(t-s)} |h'_{e^{2as}}(i)|. \end{aligned}$$

We get

Proposition 5.11. *If $r < r_c$, there is a $c < \infty$ and a subpower function φ such that if $E_t = E_{t,\beta,\psi}$ denotes the event that for $1 \leq s \leq t$,*

$$s^\beta \varphi(s)^{-1} \leq |h'_{s^2}(i)| \leq s^\beta \varphi(s), \quad (52)$$

$$(t/s)^\beta \varphi(t/s)^{-1} |h'_{s^2}(i)| \leq |h'_{t^2}(i)| \leq (t/s)^\beta \varphi(t/s) |h'_{s^2}(i)|, \quad (53)$$

then

$$\mathbb{E} [|h'_{t^2}(i)|^\lambda 1_{E_t}] \geq c.$$

Note that

$$|h'_{t^2}(i)|^\lambda 1_{E_t} \leq |h'_{s^2}(i)|^\lambda (t/s)^{\beta\lambda} \varphi(t/s)^\lambda.$$

By the definition of the event, we also get $Y_{e^{2at}} \asymp e^{at}$ and $S_{e^{2at}} \asymp 1$, so that

$$\mathbb{E} [|h'_{e^{2at}}(i)|^\lambda] \asymp [e^{at}]^{-\zeta}.$$

This is how (29) and (30) are derived.

To get (31) we have to estimate the correlations. Suppose that S, T are positive integers with $n^2/2 \leq S \leq T \leq n^2$. Let $h^{(S)}, h^{(T)}$ be defined as in the beginning of the section. Recall that $\hat{f}'_S(z) \hat{f}'_T(z)$ has the same distribution as $(h^{(S)})'(z)(h^{(T)})'(z)$. Set $h := h^{(S)}$ and $\tilde{h} := h^{(T)}$. Let $E(S), E(T)$ denote the indicator function of the event from Proposition 5.11 with $t = n$ and with h being h or $\tilde{h}$, respectively. We have already noted that $\tilde{h}_t, 0 \leq t \leq T - S$ is independent of $h_s, 0 \leq s \leq S$. Using (53) we see that

$$|\tilde{h}'_T(i)|^\lambda E(T) \leq |\tilde{h}'_{T-S}(i)|^\lambda \left(\frac{n^2}{T-S+1} \right)^{\frac{\beta\lambda}{2}} \varphi \left(\frac{n^2}{T-S+1} \right).$$

(Here we use φ for a subpower function whose exact value may change from line to line.) The random variable on the right hand side is independent of h . Therefore, using Theorem 5.4,

$$\begin{aligned} & \mathbb{E} [|h'_S(i)|^\lambda |\tilde{h}'_T(i)|^\lambda E(T)] \\ & \leq \mathbb{E} [|h'_S(i)|^\lambda] \mathbb{E} [|\tilde{h}'_{T-S}(i)|^\lambda] \left(\frac{n^2}{T-S+1} \right)^{\frac{\beta\lambda}{2}} \varphi \left(\frac{n^2}{T-S+1} \right) \\ & \leq c n^{-\zeta} (T-S+1)^{-\zeta/2} \left(\frac{n^2}{T-S+1} \right)^{\frac{\beta\lambda}{2}} \varphi \left(\frac{n^2}{T-S+1} \right) \\ & = c n^{-2\zeta} \left(\frac{n^2}{T-S+1} \right)^{\frac{\beta\lambda+\zeta}{2}} \varphi \left(\frac{n^2}{T-S+1} \right), \end{aligned}$$

and (31) follows.

REFERENCES

- [Bef08] Vincent Beffara. The dimension of the SLE curves. *Ann. Probab.*, 36(4):1421–1452, 2008.
- [JL09] Fredrik Johansson and Gregory F. Lawler. Tip multifractal spectrum for the Schramm-Loewner evolution. In preparation, 2009.
- [Law05] Gregory F. Lawler. *Conformally invariant processes in the plane*. American Mathematical Society, Providence, R.I., 2005.
- [Law09] Gregory F. Lawler. Multifractal analysis of the reverse flow for the Schramm-Loewner evolution. Preprint, 2009.
- [Lin08] Joan R. Lind. Hölder regularity of the SLE trace. *Trans. Amer. Math. Soc.*, 360(7):3557–3578, 2008.
- [LSW04] Gregory F. Lawler, Oded Schramm, and Wendelin Werner. Conformal invariance of planar loop-erased random walks and uniform spanning trees. *The Annals of Probability*, 32, 2004.
- [MR05] Donald E. Marshall and Steffen Rohde. The Loewner differential equation and slit mappings. *J. Amer. Math. Soc.*, 18(4):763–778 (electronic), 2005.
- [Pom92] Christian Pommerenke. *Boundary behaviour of conformal maps*. Springer-Vlg, Berlin, 1992.
- [RS05] Steffen Rohde and Oded Schramm. Basic properties of SLE. *Ann. of Math. (2)*, 161(2):883–924, 2005.
- [RY99] Daniel Revuz and Marc Yor. *Continuous martingales and Brownian motion*. Springer, Berlin, 3. ed. edition, 1999.
- [Sch00] Oded Schramm. Scaling limits of loop-erased random walks and uniform spanning trees. *Israel J. Math.*, 118:221–288, 2000.

FREDRIK JOHANSSON, DEPARTMENT OF MATHEMATICS, THE ROYAL INSTITUTE OF TECHNOLOGY, S – 100 44 STOCKHOLM, SWEDEN

E-mail address: frejo@math.kth.se

GREGORY F. LAWLER, DEPARTMENT OF MATHEMATICS AND DEPARTMENT OF STATISTICS, UNIVERSITY OF CHICAGO, 5734 S. UNIVERSITY AVENUE, CHICAGO, IL 60637, USA

E-mail address: lawler@math.uchicago.edu